

level of 20 milligrams of dichlorvos per kilogram of body weight for the removal of bots and ascarids. The same dosage level is repeated every 21 to 28 days for the control of bots and ascarids. For the control of bots only, the repeat dosage is 10 milligrams per kilogram of body weight every 21 to 28 days during bot fly season.

(3) Do not use this product in animals simultaneously or within a few days before or after treatment with or exposure to cholinesterase-inhibiting drugs, pesticides or chemicals. Do not administer in conjunction with or within 1 week of administration of muscle-relaxant drugs, phenothiazine derived tranquilizers, or central nervous system depressants.

(4) Do not use in horses which are severely debilitated or suffering from diarrhea or severe constipation, infectious disease, toxemia, or colic. Do not administer to horses affected with chronic alveolar emphysema (heaves) or other respiratory conditions.

(5) Do not use in horses intended for food purposes.

(6) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(i) *Conditions of use in cats and puppies.* (1) It is indicated for the removal and control of roundworms (*Toxocara canis*, *Toxocara cati*, *Toxascaris leonina*) and hookworms (*Ancylostoma caninum*, *Ancylostoma tubaeforme*, *Uncinaria stenocephala*) occurring in the intestinal tracts of cats and puppies.

(2) The drug is in tablet form and is administered orally at a dosage level of 5 mg of the active ingredient per pound of body weight.

(3) Do not administer to puppies or cats showing signs of constipation, mechanical blockage of the intestinal tract, impaired liver function, or to animals recently exposed to or showing signs of infectious disease. The drug is a cholinesterase inhibitor and should not be used simultaneously or within a few days before or after treatment with or exposure to cholinesterase-inhibiting drugs, pesticides, or chemicals.

(4) Do not use in animals under 10 days of age or under 1 pound of body weight.

(5) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[40 FR 13838, Mar. 27, 1975, as amended at 48 FR 40704, Sept. 9, 1983; 51 FR 28546, Aug. 8, 1986; 62 FR 35076, June 30, 1997]

§ 520.608 Dicloxacillin sodium monohydrate capsules.

(a) *Specifications.* Each capsule contains dicloxacillin sodium monohydrate equivalent to 50, 100, 200, or 500 milligrams of dicloxacillin.

(b) *Sponsor.* See No. 000856 in § 510.600

(c) of this chapter.

(c) *Conditions of use. Dogs—*(1) *Amount.* 5 to 10 milligrams per pound of body weight, three times daily. In severe cases, up to 25 milligrams per pound of body weight three times daily.

(2) *Indications for use.* Treatment of pyoderma (pyogenic dermatitis) due to penicillinase-producing staphylococci sensitive to the drug.

(3) *Limitations.* For the treatment of dogs only. Continue treatment for 24 to 48 hours after the animal has become afebrile or asymptomatic. Administer 1 to 2 hours before feeding to ensure maximum absorption. Not for use in animals which are raised for food production. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[57 FR 37325, Aug. 18, 1992]

§ 520.620 Diethylcarbamazine oral dosage forms.

§ 520.622 Diethylcarbamazine citrate oral dosage forms.

§ 520.622a Diethylcarbamazine citrate tablets.

(a) *Sponsors.* (1) See 015579 in § 510.600(c) of this chapter for use of 50, 200, and 400 milligram tablets for prevention of heartworm disease in dogs and as an aid in the treatment of ascarid infections in dogs and cats.

(2) See 053501 in § 510.600(c) of this chapter for use of 100, 200, and 300 milligram tablets for prevention of heartworm disease in dogs and as an aid in the treatment of ascarid infections in dogs.

(3) See 050604 in § 510.600(c) of this chapter for use of 50, 100, 200, 300, or 400